

# Ultrafast Probes for Dirac Materials

**Dmitry Yarotski**

*Center for Integrated Nanotechnologies  
Materials Physics and Applications Division  
Los Alamos National Laboratory*



*Quantum and Dirac Materials Workshop  
March 8-11, 2015, Santa Fe, NM, USA*



# Collaborators and Acknowledgements

**LANL Staff:** Rohit Prasankumar, Antoinette Taylor, Abul Azad, Steve Gilbertson, George Rodriguez, Tomasz Durakiewicz, Aditya Mohite, Andrew Dattelbaum, Quanxi Jia, Stuart Trugman, Jian-xin Zhu

**LANL Postdocs:** Rolando Valdes Aguilar, Yaomin Dai, Keshav Dani, John Bowlan, Jingbo Qi, Jinho Lee, Georgi Dakovski

**Brookhaven National Laboratory:** Genda Gu, Ruidan Zhong

**Rutgers University:** Matthew Brahlek, Namrata Bansal, Seongshik Oh



**Rice University:** Sina Najmaei, Jun Lou, Pulickel M. Ajayan,

We gratefully acknowledge the support from the U.S. Department of Energy through the Center for Integrated Nanotechnologies, LANL LDRD Program, and the UC Office of the President under UC Lab Fees Research Program



# Why Ultrafast Spectroscopy ?





Ultrafast (10-100 fs) spectroscopy can resolve non-equilibrium dynamics (quasiparticle, transport etc.) at the fundamental time and spatial scales of electronic and nuclear motion





# Ultrafast Coherent Order Manipulation


Manipulation of order parameters ♦ Photoinduced phase transitions ♦  
New non-thermally accessible phases.








# Graphene: The Slice that Started It All




- **Graphene:** a basis for 0D buckyballs, 1D carbon nanotubes, and 3D graphite
- Quasiparticles are described by relativistic Dirac equation – **Dirac Material**
- Massless Dirac quasiparticles exhibit novel transport properties (high mobility, excellent conductivity)

Understanding the *non-equilibrium behavior of photoexcited graphene* is important for science and applications in detectors, solar cells and displays.



Bae et al. *Nat. Nanotech.* 2010



Bonaccorso et al. *Nat. Photonics* 2010



# Quasiparticles in Graphene

Linear dispersion near Dirac point gives for relativistic quasiparticles:



$$E \approx \hbar v_F k$$

$$E_F^{e,h} \sim \hbar v_F \sqrt{\pi N_{e,h}}$$

**Are photoexcited quasiparticles in graphene relativistic too?**



**Two types of optical conductivity in graphene:**

**Interband** is *constant* in a wide spectral range (flat 2.3% absorption)



Mak et al., Phys. Rev. Lett. (2008)

**Intraband** differs for linear and parabolic bands



$$\sigma_{intra}^{parab} = \frac{Ne^2}{m} \frac{\gamma}{\omega^2 + \gamma^2}$$

$$\sigma_{intra} \propto \sqrt{N} \frac{\gamma}{\omega^2 + \gamma^2}$$



**Measuring conductivity change after photoexcitation as function of  $N$  will indicate whether non-equilibrium quasiparticles are relativistic**



# Measuring Relativistic Quasiparticles in Graphene

We measure the photoinduced conductivity change:


$$\Delta\sigma = (\sigma_{inter} + \sigma_{intra}) \Big|_{Photo-excited} - (\sigma_{inter} + \sigma_{intra}) \Big|_{Intrinsic\ doping}$$



The change in conductivity, as measured in a visible pump-probe experiment, is dominated by the intraband component!



# Near-IR Pump, Visible-Probe Spectroscopy



- ❖ 1.55 eV pump, 1.77 eV probe experiments
- ❖ Fermi energy after photoexcitation = 700 meV (for  $N \sim 3.1 \times 10^{13} / \text{cm}^2$ )
- ❖ Decay dynamics are qualitatively identical for all photon energies (1.74-2.42 eV)
- ❖ Electron-electron thermalization within <100 fs – **Amplitude gives optical  $\Delta\sigma$**
- ❖ Electron-phonon thermalization within 1.4 ps






# Hot Dirac Fermions in Graphene

Intraband contribution

Interband contribution




Reflectivity (or conductivity) change follows  $\sqrt{N}$  from  $E_F^{e,h} \sim \hbar v_F \sqrt{\pi N_{e,h}}$

**Our experiment reveals the relativistic nature of photoexcited Dirac quasiparticles in graphene**





# Time-Resolved ARPES

## High Harmonic Generation – Extreme nonlinear frequency upconversion



M. Ferray, *et al.* J. Phys., 21 (1988); P.B. Corkum, PRL 71, 1994 (1993)




### STATIC ARPES:

- ◆ probes electronic structure in both **E** and **k** domains


### DYNAMIC ARPES:

- ◆ probes **transient** electronic structure changes in both **E** and **k** domains
- ◆ **Fills** excited states to **reveal** their structure








# Photoexcited Fermi-Dirac Distribution in Graphene



- ❖ Is the Fermi-Dirac distribution of photoexcited carriers in graphene more like a **metal** (same  $\mu_e$  and  $\mu_h$ ) or like a **semiconductor** (separate  $\mu_e$  and  $\mu_h$ )?
- ❖ Do processes like Auger recombination influence the dynamics at early times?
- ❖ Time-resolved photoemission experiments show that, in our samples, the photoexcited carriers retain separate F-D distributions for a few hundred femtoseconds




# Recombination of Electronic States in Graphene




- ❖ Ultrafast pump/probe experiment on CVD grown graphene
  - 30 fs IR pump and sub-10 fs, 30-eV probe via HHG
  - measure tr-ARPES
- ❖ A short-lived distribution of carriers and holes is formed after optical excitation.
- ❖ Separate populations are:
  - ◆ semi-conductor like ( $\mu^* \neq 0$ ) at early delays
  - ◆ metallic like ( $T^* \neq 0$ ) at later times



# Topological Insulators




\* after A. Lanzara



## Materials with exotic surface states

- Linear  $E$ - $k$  dispersion
- TRS protection against scattering
- Locked spin- $k$  relationship
- Majorana Fermions
- Spintronics, optoelectronics



- Real materials are not ideal – dopants/defects result in significant bulk interference
- THz spectroscopy provides the ability to separate the collective motion of charge carriers in bulk vs. surface states








# Optical Pump Terahertz Probe




↓  
① Fourier Transform




Complex Transmissivity & Fresnel Analysis

↓  
②

$$T(\omega) = \frac{E(\omega)_{\text{sample}}}{E(\omega)_{\text{ref}}} = \frac{1+n_3}{1+n_3+\sigma(\omega)dZ_o}$$




③ Extract Complex Conductivity






# Terahertz Conductivity of $\text{Bi}_2\text{Se}_3$





- **Low freq. spectra:**  
Drude component:  $1/\tau \sim 1$  THz  
Bulk phonon:  $\omega_0 \sim 1.9$  THz
- **Electron density consistent with**  
 $n_{\text{surf}} \sim 1.5 \times 10^{13} \text{ cm}^{-2}$
- **Drude term is thickness independent**  
Surface.
- **Phonon is not  $\rightarrow$  Bulk effect.**





# Time-Resolved THz Spectroscopy

Fix THz gate delay at maximum and scan pump-probe delay





# Photo-Induced Conductivity in $\text{Bi}_2\text{Se}_3$

## ➤ Drude-Lorentz Model:



$$G_j(\omega) = \left( -\frac{\omega_{\text{pD},j}^2}{i\omega - \Gamma_{\text{Drude},j}} - \frac{i\omega\omega_{\text{pDL},j}^2}{\omega_{\text{DL},j}^2 - \omega^2 - i\omega\Gamma_{\text{Lorentz},j}} - i(\epsilon_\infty - 1)\omega \right) \epsilon_0 d$$

## ➤ Well described by *single* carrier type

## ➤ Carriers in 20 QL decay faster



## ➤ **Green**: Drude (free electron).

## ➤ **Purple**: Phonon.






# Photo-Induced Drude Properties in 20 QL



**Low Fluence:** increase scat. rate  $\rightarrow$  increase T


**High Fluence:** increase plasma freq.  $\rightarrow$  decrease T









# Photo-Induced Phonon Frequency Shift in 20 QL




- At high fluence, phonon shifts - similar to increase in temperature.
- Highest lattice temperature ~ 200 K



# Photo-Induced Drude Properties in 10 QL



- Plasma frequency doesn't change as much as in 20 QL sample.
- Scattering rate does, so the sample becomes more transparent at higher fluence.






# Physical Picture

Phonon-induced bulk-to-surface scattering is not effective below  $T_D=180\text{ K}$

Wang et al., Phys. Rev. Lett. 109, 127401 (2012)  
Sim et al., Phys. Rev. B 89, 165137 (2014)



**Hot surface carriers can be accessed independently from the bulk ones using THz spectroscopy**

**Thin 10 QL films are similar to graphene:**

- ❖ Surface electrons dominate, but  $\Delta\omega_p$  is small
- ❖  $\Gamma_{\text{surf}}$  increases due to e-h scattering and temperature rise ( $\sim 200\text{ K}$ ) due to e-ph relaxation

**Thick 20 QL films:**

- ❖ Surface response dominates at low fluences
- ❖ High fluences result in large number of bulk carriers  $\Rightarrow$  higher  $\Delta\omega_p$  and  $\Gamma_{\text{bulk}}$
- ❖ Bulk electrons decay in  $\sim 5\text{ ps}$
- ❖ Surface electrons decay in  $20\text{ ps}$  preserving high scattering rates



# Topological Crystalline Insulators

PRL **106**, 106802 (2011)

PHYSICAL REVIEW LETTERS

week ending  
11 MARCH 2011

## Topological Crystalline Insulators

Liang Fu

*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*


(Received 5 October 2010; revised manuscript received 31 December 2010; published 8 March 2011)

TI  $\longrightarrow$  Time Reversal Symmetry

TCI  $\longrightarrow$  Crystalline Symmetry


Metallic states on **High Symmetry** surfaces!

(001) surface






# Topological Phase Transition in $\text{Pb}_{1-x}\text{Sn}_x\text{Se}$



$\text{Pb}_{0.77}\text{Sn}_{0.23}\text{Se}$



Dziawa et al. Nat. Mater. **11**, 1023 (2012)

**P**-induced TPT in  $\text{Pb}_{1-x}\text{Sn}_x\text{Se}$

Xi et al. PRL **113**, 096401 (2014)





# Topological Phase Transition in $\text{Pb}_{1-x}\text{Sn}_x\text{Te}$


**PbTe**  
Trivial

Doping-driven Topological phase transition



$\text{Pb}_{1-x}\text{Sn}_x\text{Te}$

**SnTe**  
TCI



**$X_c = 0.4$  at 5K**


Yan et al. PRL **112**, 186801 (2014)

Can we use UOS to find the evidence for TPT with  
*temperature* and *doping*?




# Preliminary Results and Future Directions

## Doping-induced TPT at 5 K



## Temperature-induced TPT at $x=0.4$



- Strong electron-phonon coupling in TI state – common to all TI
- Investigate the effect of magnetic field using THz spectroscopy to probe conductivity of photoexcited carriers.
- Apply circularly polarized pump to break TRS and study the dynamics of the  $\mathbf{k}$ -spin locking process.



# Temperature Dependence of Decay Amplitudes

